



PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM	Bo	
Search	PubMed	▼	for					Go	Clear
		Limits	Preview/Index	History	Clipboard	Details			
Display		Abstract	▼	Sort	▼	Save	Text	Clip Add	Order

Entrez PubMed

☐ 1: Cell 1990 Feb 9;60(3):461-72

Related Articles, Domains, Nucleotide, Protein, Books, LinkOut

**A novel octamer binding transcription factor is differentially expressed in mouse embryonic cells.**

PubMed Services

**Okamoto K, Okazawa H, Okuda A, Sakai M, Muramatsu M, Hamada H.**

Department of Biochemistry, Faculty of Medicine, University of Tokyo, Japan.

Related Resources

We have identified a novel octamer binding factor (Oct-3) in P19 embryonal carcinoma cells. Oct-3, which recognizes the typical octamer motif (ATTTGCAT) as well as the AT-rich sequence TTAAAATTCA, is present in P19 stem cells but disappears when the cells are induced to differentiate by retinoic acid (RA). Cloned cDNA corresponding to Oct-3 encodes a protein of 377 amino acids. Oct-3 has a conserved POU domain, but the remaining part is distinct from other POU domain-containing proteins such as Oct-1 and Oct-2. mRNA of 1.5 kb coding for Oct-3 is abundant in P19 stem cells but is dramatically repressed during RA-induced differentiation. Repression of the 1.5 kb mRNA is rapid and specific to RA. In mouse, oct-3 mRNA is undetectable in all the adult organs examined. The N-terminal proline-rich region of Oct-3, when fused to the DNA binding domain of c-Jun, functions as a transcriptional activating domain. We suggest that Oct-3 is a novel octamer binding transcription factor that is developmentally regulated during mouse embryogenesis.

PMID: 1967980 [PubMed - indexed for MEDLINE]

Display	Abstract	▼	Sort	▼	Save	Text	Clip Add	Order
---------	----------	---	------	---	------	------	----------	-------

Write to the Help DeskNCBI | NLM | NIHDepartment of Health & Human ServicesFreedom of Information Act | Disclaimer



PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM	Bo	
Search	PubMed	▼	for					Go	Clear
		Limits	Preview/Index	History	Clipboard	Details			
Display		Abstract	▼	Sort	▼	Save	Text	Clip Add	Order

Entrez PubMed

☐ 1: Nature 1990 Jun 21;345(6277):686-92 Related Articles, Protein, Books, LinkOut

### **A POU-domain transcription factor in early stem cells and germ cells of the mammalian embryo.**

PubMed Services

**Rosner MH, Vigano MA, Ozato K, Timmons PM, Poirier F, Rigby PW, Staudt LM.**

Metabolism Branch, National Cancer Institute, Bethesda, Maryland 20892.

Related Resources

The murine oct-3 gene encodes a transcription factor containing a POU-specific domain and a homeodomain. In marked contrast to other homeodomain-encoding genes, oct-3 is expressed in the totipotent and pluripotent stem cells of the pregastrulation embryo and is down-regulated during differentiation to endoderm and mesoderm, suggesting that it has a role in early development. The oct-3 gene is also expressed in primordial germ cells and in the female germ line.

PMID: 1972777 [PubMed - indexed for MEDLINE]

Display	Abstract	▼	Sort	▼	Save	Text	Clip Add	Order
---------	----------	---	------	---	------	------	----------	-------

Write to the Help Desk  
NCBI | NLM | NIH  
Department of Health & Human Services  
Freedom of Information Act | Disclaimer

L Number	Hits	Search Text	DB	Time stamp
1	23	cicm	USPAT; EPO; DERWENT	2001/12/12 07:16
2	3	cicm with bovine	USPAT; EPO; DERWENT	2001/12/12 07:16
3	163	inner adj1 cell adj1 mass	USPAT; EPO; DERWENT	2001/12/12 07:17
4	24	(inner adj1 cell adj1 mass) same (bovine or cow)	USPAT; EPO; DERWENT	2001/12/12 07:18
5	24	((inner adj1 cell adj1 mass) or cicm) same (bovine or cow)	USPAT; EPO; DERWENT	2001/12/12 07:18